

Low-Energy Spin Dynamics of CuO Chains in $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$

D.N. Aristov ^{a,1}, D.R. Grempel ^{b,2}

^a*Petersburg Nuclear Physics Institute, Gatchina 188350 St.Petersburg, Russia*

^b*Service de Physique de l'État Condensé, CEA-Saclay, F-91191 Gif-sur-Yvette, France*

Abstract

We study the spin fluctuation dynamics of Cu-O chains in the oxygen deficient planes of $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$. The chains are described by a model including antiferromagnetic interactions between the spins and Kondo-like scattering of the oxygen holes by the copper spins. There are incommensurate spin fluctuations along the direction of the chains. The dynamic structure factor of this system is qualitatively different from that of a quasi one-dimensional localized antiferromagnet due to the presence of itinerant holes. We compute the dynamic structure factor that could be measured in neutron scattering experiments.

Key words: Quantized spin models; Dynamic properties; Y-based cuprates; Neutron scattering techniques

The oxygen deficient CuO_x planes play an important role in the physics of the high- T_c compounds $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$. These planes are composed of $\text{Cu}_{n+1}\text{O}_n$ chain fragments in which Cu and O atoms form a linear alternating chain. The physical properties of the chain planes are dominated by the longest chain fragments. To a first approximation, these chains may be viewed as decoupled from each other and from the CuO_2 planes. Therefore, important one-dimensional fluctuation effects are expected in the charge and spin excitation spectra of the CuO_x planes.

The low-energy spin dynamics of this system was studied theoretically in [1] within a strong-coupling model in which every Cu site is occupied by a hole

while the average filling of the O sites $c < 1$. Holes can hop between the O sites obeying a constraint of no double occupancy. There is an exchange interaction between the spins of all pairs of adjacent holes. In this model, a spin *sequence* on the chain is not affected by the oxygen-hole motion, *i.e.*, there is spin-charge separation. Due to this fact, the energy spectrum of the system can be computed in closed form. The spin-spin correlation function of the chains is non-trivial, however. It is given by a convolution of two functions. The first one is the dynamic structure factor of a one-dimensional Heisenberg antiferromagnet on a “fictitious” lattice containing a number of sites equal to the total number of holes in the chain. The second factor is related to the density-density correlation function of the itinerant oxygen holes and describes the evolution of the spin *positions* as determined by the motion of the holes. The energy scales character-

¹ Corresponding author. E-mail: aristov@thd.pnpi.spb.ru

² On leave from Service de Physique Statistique, Magnétisme et Supraconductivité, CEA-Grenoble, F-38054 Grenoble, France.

izing these two factors are the spin-wave velocity v_S and the Fermi velocity v_F , respectively. For $c \simeq 0.3$, the parameterization of [1] gives $v_S = 150$ meV and $v_F = 800$ meV *i.e.* $v_S \ll v_F$.

An appropriate tool for the experimental study of the magnetic fluctuations is inelastic scattering of spin polarized neutrons. The magnetic intensity is modulated along the direction of the chains and peaks at the incommensurate position, $Q_{\max} = \pi(1 + c)$. The spectrum of magnetic excitations at $q = Q_{\max}$ extends up to the energies of order of the Fermi energy of the itinerant holes. The absolute value of the neutron cross-section at $Q_{\max}$ was estimated in [1]. Here we compute the shape of the magnetic signal near the maximum as predicted by our model.

Up to fundamental constants, the neutron's differential cross-section for magnetic scattering is given by the expression :

$$\frac{d\sigma(q, \omega)}{d\Omega_s d\omega} \propto \sqrt{\frac{E_i - \omega}{E_i}} S(q, \omega), \quad (1)$$

with E_i the energy of the incident neutron, q the momentum transfer in the direction parallel to the chain and ω is the neutron's energy loss. The form of the spin-spin correlation function $S(q, \omega)$ was found in [1]. Near $q = Q_{\max}$ we may keep only the principal term and linearize the dispersion relation of both spin and density fluctuations. Up to a coefficient we have

$$S(Q_{\max} + q, \omega) \simeq \int_0^\omega d\varepsilon \int_{-\infty}^{\infty} \frac{dk}{\sqrt{v_F |k|}} \frac{1}{[\varepsilon^2 - v_F^2 k^2]^{1/4}} \times [(\omega - \varepsilon)^2 - v_S^2 (q - k)^2]^{-1/2}. \quad (2)$$

The shape of the function $S(q, \omega)$ is shown in the upper half of the Fig. 1.

Consider an experimental geometry in which the momentum of the outgoing neutron is in the direction perpendicular to chains and all the neutrons are collected, irrespective to their final energy. The measured intensity is proportional to the integral $I(q) = \int_0^{E_i} d\omega \frac{d\sigma(q, \omega)}{d\Omega_s d\omega}$. Experimentally, one typically uses polarized neutrons with incident en-




Fig. 1. The dynamic structure factor of the chains $S(q, \omega)$ is shown in the upper panel. The predicted integrated signal for an incident neutron energy $E_i = 25$ meV is plotted in the lower part of the figure.

ergy $E_i = 25$ meV. We show the corresponding curve $I(q)$ in the lower half of the Fig. 1.

In summary, we calculated the shape of a magnetic signal from the chains in $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$, visible in neutron experiments. Albeit weak, this signal could be observed in a high flux reactor.

One of us (D.A.) acknowledges partial financial support from the Russian State Program for Statistical Physics (Grant VIII-2) and the Russian Program "Neutron Studies of Condensed Matter".

References

- [1] D.N. Aristov and D. R. Grempel, Phys. Rev. B **55**, 11358, (1997) ; D.N. Aristov, *ibid.* **57**, 12825, (1998).